

On the mean square of the error term for the two-dimensional divisor problem (I)

Wenguang Zhai¹, Xiaodong Cao²

¹ Shandong Normal University, Jinan, P. R. China

² Beijing Institute of Petro-Chemical Technology, Beijing, P. R. China

Abstract. Suppose a and b are two fixed positive integers such that $(a, b) = 1$. In this paper we shall establish an asymptotic formula for the mean square of the error term $\Delta_{a,b}(x)$ of the general two-dimensional divisor problem.

1 Introduction

Suppose $1 \leq a \leq b$ are two fixed integers. Without loss of generality, we suppose $(a, b) = 1$. Define $d_{a,b}(n) := \sum_{n=ha_r b} 1$. The general two-dimensional divisor problem is to study the error term

$$\Delta_{a,b}(x) := \sum_{n \leq x} d_{a,b}(n) - \zeta(b/a)x^{1/a} - \zeta(a/b)x^{1/b},$$

if $a \neq b$. If $a = b$, then the appropriate limit is to be taken in the above sum. This problem attracts the interests of many authors.

When $a = b = 1$, $\Delta_{1,1}(x)$ is the error term of the well-known Dirichlet divisor problem. Dirichlet first proved that $\Delta_{1,1}(x) = O(x^{1/2})$. The exponent $1/2$ was improved by many authors. The latest result reads(see Huxley[9])

$$(1.1) \quad \Delta_{1,1}(x) \ll x^{131/416}(\log x)^{26947/8320}.$$

For the lower bounds, the best results read

$$(1.2) \quad \Delta_{1,1}(x) = \Omega_+(x^{1/4}(\log x)^{1/4}(\log \log x)^{(3+\log 4)/4} \exp(-c\sqrt{\log \log \log x}))(c > 0)$$

2000 Mathematics Subject Classification: 11N37.

Key Words: two-dimensional divisor problem, error term, mean square, asymptotic formula .

This work is supported by National Natural Science Foundation of China(Grant No. 10771127) and National Natural Science Foundation of Shandong Province(Grant No. Y2006A31).

and

$$(1.3) \quad \Delta_{1,1}(x) = \Omega_-(x^{1/4} \exp(c'(\log \log x)^{1/4}(\log \log \log x)^{-3/4})) \quad (c' > 0),$$

which are due to Hafner[5] and Corrádi and Kátai[2], respectively.

When $a \neq b$, Richert[20] proved that

$$(1.4) \quad \Delta_{a,b}(x) \ll \begin{cases} x^{\frac{2}{3(a+b)}}, & b \leq 2a, \\ x^{\frac{2}{5a+2b}}, & b \geq 2a. \end{cases}$$

Better upper estimates can be found in [14, 15, 19, 21]. Hafner[6] proved that

$$(1.5) \quad \Delta_{a,b}(x) = \Omega_+(x^{1/2(a+b)}(\log x)^{b/2(a+b)} \log \log x)$$

and

$$(1.6) \quad \Delta_{a,b}(x) = \Omega_-(x^{1/2(a+b)} e^{U(x)}),$$

where

$$U(x) = B(\log \log x)^{b/2(a+b)}(\log \log \log x)^{b/2(a+b)-1}$$

for some positive constant $B > 0$.

It is conjectured that the estimate

$$(1.7) \quad \Delta_{a,b}(x) = O(x^{1/2(a+b)+\varepsilon})$$

holds for any $1 \leq a \leq b$, $(a, b) = 1$. When $a = b = 1$, the conjecture (1.7) is supported by the power moment results of $\Delta_{1,1}(x)$. For the mean square of $\Delta_{1,1}(x)$, Cramér[3] first proved the classical result

$$(1.8) \quad \int_1^T \Delta_{1,1}^2(x) dx = \frac{(\zeta(3/2))^4}{6\pi^2\zeta(3)} T^{3/2} + O(T^{5/4+\varepsilon}).$$

The estimate $O(T^{5/4+\varepsilon})$ in (1.8) was improved to $O(T \log^5 T)$ in [22] and $O(T \log^4 T)$ in [18]. The higher-power moments of $\Delta_{1,1}(x)$ were studied in [7, 13, 23, 25, 26, 27]. When $a \neq b$, by using the theory of the Riemann zeta-function, Ivić[12] proved that

$$(1.9) \quad \int_1^T \Delta_{a,b}^2(x) dx \begin{cases} \ll T^{1+1/(a+b)} \log^2 T, \\ = \Omega(T^{1+1/(a+b)}). \end{cases}$$

He also conjectured that the asymptotic formula

$$(1.10) \quad \int_1^T \Delta_{a,b}^2(x) dx = c_{a,b} T^{1+1/(a+b)} (1 + o(1))$$

holds for some positive constant $c_{a,b}$. We note that the Ω -result in (1.9) is also contained in a very general result of the second-named author (see Theorem 5 of [1]).

When $a = b = 1$, the proofs of most power moment results of $\Delta_{1,1}(x)$ mentioned above were started from the well-known truncated Voronoi's formula (see, for example, [10])

$$(1.11) \quad \Delta_{1,1}(x) = (\sqrt{2}\pi)^{-1} x^{1/4} \sum_{n \leq N} \frac{d(n)}{n^{3/4}} \cos(4\pi\sqrt{xn} - \frac{\pi}{4}) + O(x^\varepsilon + x^{1/2+\varepsilon} N^{-1/2})$$

for $1 \ll N \ll x^A$, where $A > 0$ is any fixed constant. Note that the infinite series $\sum_{n=1}^{\infty} d(n)n^{-3/4} \cos(4\pi\sqrt{xn} - \pi/4)$ is conditionally convergent. When $a \neq b$, Krätzel[14] provided a series representation of $\Delta_{a,b}(x)$. However, Krätzel's series converges only when $b < 3a/2$, which is a significant restriction for many applications.

The aim of this paper is to prove Ivić's conjecture (1.10). More precisely, we shall prove the following Theorem.

Theorem . Suppose $1 \leq a < b$ are fixed integers for which $(a, b) = 1$. Then we have

$$(1.12) \quad \int_1^T \Delta_{a,b}^2(x) dx = c_{a,b} T^{\frac{1+a+b}{a+b}} + O(T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \log^{7/2} T),$$

where

$$c_{a,b} := \frac{a^{b/(a+b)} b^{a/(a+b)}}{2(a+b+1)\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n),$$

$$g_{a,b}(n) := \sum_{n=h^a r^b} h^{-\frac{a+2b}{2a+2b}} r^{-\frac{b+2a}{2a+2b}}.$$

Remark 1. It is easy to see that the function $g_{a,b}(n)$ is symmetric for a and b , namely $g_{a,b}(n) = g_{b,a}(n)$. The convergence of the infinite series $\sum_{n=1}^{\infty} g_{a,b}^2(n)$ will be proved in Section 4.

Remark 2. Our theorem also holds for $a = b = 1$. In this case we have

$$g_{1,1}(n) = d(n)n^{-3/4}, \quad c_{1,1} = \frac{1}{6\pi^2} \sum_{n=1}^{\infty} d^2(n)n^{-3/2} = \frac{(\zeta(3/2))^4}{6\pi^2\zeta(3)},$$

where $d(n) = d_{1,1}(n)$ is the Dirichlet divisor function. Hence our Theorem provides a new proof of Cramer's classical result (1.8).

Notations. $\mathbb{Z}$ denotes the set of all integers. For a real number u , $[u]$ denotes the integer part of u , $\{u\}$ denotes the fractional part of u , $\psi(u) = \{u\} - 1/2$, $\|u\|$ denotes the distance from u to the integer nearest to u . $\mu(n)$ is the Möbius function,

(m, n) denotes the greatest common divisor of natural numbers m and n . $n \sim N$ means $N < n \leq 2N$. ε always denotes a sufficiently small positive constant. $SC(\Sigma)$ denotes the summation condition of the sum Σ when it is complicated. Finally, define

$$\sum'_{\alpha \leq n \leq \beta} f(n) = \begin{cases} \sum_{\alpha < n < \beta} f(n), & \alpha \notin \mathbb{Z}, \beta \notin \mathbb{Z}, \\ f(\alpha)/2 + \sum_{\alpha < n < \beta} f(n), & \alpha \in \mathbb{Z}, \beta \notin \mathbb{Z}, \\ f(\alpha)/2 + \sum_{\alpha < n < \beta} f(n) + f(\beta)/2, & \alpha \in \mathbb{Z}, \beta \in \mathbb{Z}. \end{cases}$$

2 Two preliminary Lemmas

In order to prove our theorem, we need the following two Lemmas. Lemma 2.1 is well-known; see for example, Heath-Brown[8]. Lemma 2.2 is Theorem 2.2 of Min[17], see also Lemma 6 of Chapter 1 in [24]. A weaker version of Lemma 2.2 can be found in [16], which also suffices for our proof.

Lemma 2.1. Let $H \geq 2$ be any real number. Then

$$\psi(u) = - \sum_{1 \leq |h| \leq H} \frac{e(hu)}{2\pi i h} + O\left(\min(1, \frac{1}{H\|u\|})\right).$$

Lemma 2.2. Suppose $A_1, \dots, A_5$ are absolute positive constants, $f(x)$ and $g(x)$ are algebraic functions in $[a, b]$ and

$$\begin{aligned} \frac{A_1}{R} &\leq |f''(x)| \leq \frac{A_2}{R}, \quad |f'''(x)| \leq \frac{A_3}{RU}, \quad U \geq 1, \\ |g(x)| &\leq A_4 G, \quad |g'(x)| \leq A_5 G U_1^{-1}, \quad U_1 \geq 1, \end{aligned}$$

$[\alpha, \beta]$ is the image of $[a, b]$ under the mapping $y = f'(x)$, then

$$\begin{aligned} \sum_{a < n \leq b} g(n) e(f(n)) &= e^{\pi i/4} \sum_{\alpha \leq u \leq \beta} b_u \frac{g(n_u)}{\sqrt{f''(n_u)}} e(f(n_u) - u n_u) \\ &\quad + O(G \log(\beta - \alpha + 2) + G(b - a + R)(U^{-1} + U_1^{-1})) \\ &\quad + O\left(G \min\left[\sqrt{R}, \max\left(\frac{1}{<\alpha>}, \frac{1}{<\beta>}\right)\right]\right), \end{aligned}$$

where n_u is the solution of $f'(n) = u$,

$$< t > = \begin{cases} \|t\|, & \text{if } t \text{ not an integer,} \\ \beta - \alpha, & \text{if } t \text{ an integer,} \end{cases}$$

$$b_u = \begin{cases} 1, & \text{if } \alpha < u < \beta, \text{ or } \alpha, \beta \text{ not integers,} \\ 1/2, & \text{if } \alpha \text{ or } \beta \text{ are integers,} \end{cases}$$

$$\sqrt{f''} = \begin{cases} \sqrt{f''}, & \text{if } f'' > 0, \\ i\sqrt{|f''|}, & \text{if } f'' < 0. \end{cases}$$

3 A Voronoi type formula of $\Delta(a, b; x)$

It suffices for us to evaluate the integral $\int_T^{2T} \Delta_{a,b}^2(x) dx$, where $T \geq 10$ is a large parameter.

It is well-known that (see for example, Ivić[11], eq.(14.46))

$$(3.1) \quad \Delta_{a,b}(x) = f(a, b; x) + f(b, a; x) + O(1),$$

where

$$f(a, b; x) := - \sum_{m \leq x^{1/(a+b)}} \psi \left(\frac{x^{1/a}}{m^{b/a}} \right).$$

Suppose $T \leq x \leq 2T$, H is a parameter such that $T^\varepsilon \ll H \ll T^{100(a+b)}$. By Lemma 2.1 we have

$$(3.2) \quad \begin{aligned} f(a, b; x) &= R_1(a, b; x) + R_2(a, b; x), \\ R_1(a, b; x) : &= \frac{1}{2\pi i} \sum_{1 \leq |h| \leq H} \frac{1}{h} \sum_{m \leq x^{1/(a+b)}} e \left(\frac{hx^{1/a}}{m^{b/a}} \right), \\ R_2(a, b; x) : &= O \left(\sum_{m \leq x^{1/(a+b)}} \min \left(1, \frac{1}{H \left\| \frac{x^{1/a}}{m^{b/a}} \right\|} \right) \right). \end{aligned}$$

Define

$$\begin{aligned} c &:= (2ab)^{ab}, J := [(\mathcal{L}/(a+b) - \log \mathcal{L}) \log^{-1} c], \mathcal{L} := \log T, \\ m_j &:= x^{1/(a+b)} c^{-j} \quad (j \geq 0). \end{aligned}$$

It is easy to see that

$$c^J \asymp T^{1/(a+b)} \mathcal{L}^{-1}.$$

We have

$$\begin{aligned}
(3.3) \quad R_1(a, b; x) &= \frac{1}{2\pi i} \sum_{1 \leq |h| \leq H} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(\frac{hx^{1/a}}{m^{b/a}}\right) + O(\mathcal{L}^2) \\
&= \frac{1}{2\pi i} \sum_{-H \leq h \leq -1} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(\frac{hx^{1/a}}{m^{b/a}}\right) \\
&\quad + \frac{1}{2\pi i} \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(\frac{hx^{1/a}}{m^{b/a}}\right) + O(\mathcal{L}^2) \\
&= -\frac{1}{2\pi i} \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(-\frac{hx^{1/a}}{m^{b/a}}\right) \\
&\quad + \frac{1}{2\pi i} \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(\frac{hx^{1/a}}{m^{b/a}}\right) + O(\mathcal{L}^2) \\
&= -\frac{\Sigma_1}{2\pi i} + \frac{\overline{\Sigma_1}}{2\pi i} + O(\mathcal{L}^2),
\end{aligned}$$

say, where

$$\Sigma_1 = \sum_{1 \leq h \leq H} \frac{1}{h} \sum_{j=0}^J \sum_{m_{j+1} < m \leq m_j} e\left(-\frac{hx^{1/a}}{m^{b/a}}\right).$$

Let

$$S_{h,j}(x) := \sum_{m_{j+1} < m \leq m_j} e\left(-\frac{hx^{1/a}}{m^{b/a}}\right).$$

Define

$$\begin{aligned}
c_1(a, b) &:= a^{b/2(a+b)} b^{a/2(a+b)} (a+b)^{-1/2}, \\
c_2(a, b) &:= a^{b/(a+b)} b^{-b/(a+b)} + b^{a/(a+b)} a^{-a/(a+b)}.
\end{aligned}$$

It is easy to check that

$$(3.4) \quad c_1(a, b) = c_1(b, a), \quad c_2(a, b) = c_2(b, a).$$

By Lemma 2.2 we get

$$\begin{aligned}
(3.5) \quad S_{h,j}(x) &= c_1(a, b) x^{\frac{1}{2(a+b)}} \sum'_{n_{j,h}(a,b) \leq r \leq n_{j+1,h}(a,b)} h^{\frac{a}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} \\
&\quad \times e\left(-c_2(a, b) x^{\frac{1}{a+b}} (h^a r^b)^{\frac{1}{a+b}} - \frac{1}{8}\right) + O(\mathcal{L}),
\end{aligned}$$

where

$$n_{j,h}(a,b) := \frac{b}{a} h(2ab)^{(a+b)bj}.$$

Inserting (3.5) into Σ_1 we get

$$(3.6) \quad \begin{aligned} \Sigma_1 &= c_1(a,b) x^{\frac{1}{2(a+b)}} \sum_{1 \leq h \leq H} \sum'_{\frac{bh}{a} \leq r \leq n_{J+1,h}(a,b)} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} \\ &\times e \left(-c_2(a,b) x^{\frac{1}{a+b}} (h^a r^b)^{\frac{1}{a+b}} - \frac{1}{8} \right) + O(\mathcal{L}^2). \end{aligned}$$

From (3.3) and (3.6) we get

$$(3.7) \quad R_1(a,b;x) = R_1^*(a,b;x) + O(\mathcal{L}^2),$$

where

$$\begin{aligned} R_1^*(a,b;x) : &= \frac{c_1(a,b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq h \leq H} \sum'_{\frac{bh}{a} \leq r \leq n_{J+1,h}(a,b)} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} \\ &\times \cos \left(2\pi c_2(a,b) x^{\frac{1}{a+b}} (h^a r^b)^{\frac{1}{a+b}} - \frac{\pi}{4} \right). \end{aligned}$$

Define

$$\begin{aligned} g(a,b;n,H,J) : &= \sum'_{\substack{n=h^a r^b, 1 \leq h \leq H \\ bh/a \leq r \leq n_{J+1,h}(a,b)}} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}}, \\ g(a,b;n) : &= \sum'_{\substack{n=h^a r^b \\ bh/a \leq r}} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}}. \end{aligned}$$

It is easy to check that if $h^a r^b \leq \min(H^{a+b}, T^{b/a}) \mathcal{L}^{-b-b^2/a-1}$, $bh/a \leq r$, then it follows that $h \leq H, r \leq n_{J+1,h}(a,b)$. Thus

$$(3.8) \quad g(a,b;n,H,J) = g(a,b;n), \quad n \leq \min(H^{a+b}, T^{b/a}) \mathcal{L}^{-b-b^2/a-1}.$$

So we have that

$$\begin{aligned} R_1^*(a,b;x) &= \frac{c_1(a,b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq H^a n_{J+1,H}(a,b)} g(a,b;n,H,J) \\ &\times \cos \left(2\pi c_2(a,b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right) \end{aligned}$$

Similarly we have

$$(3.9) \quad R_1(b, a; x) = R_1^*(b, a; x) + O(\mathcal{L}^2),$$

where

$$\begin{aligned} R_1^*(b, a; x) &= \frac{c_1(b, a)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq H^b n_{J+1, H}(b, a)} g^*(b, a; n, H, J) \\ &\quad \times \cos \left(2\pi c_2(b, a) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right), \\ g(b, a; n, H, J) : &= \sum'_{\substack{n = h^b r^a, 1 \leq h \leq H \\ ah/b \leq r \leq n_{J+1, h}(b, a)}} h^{-\frac{b+2a}{2(a+b)}} r^{-\frac{2b+a}{2(a+b)}}, \\ g(b, a; n) : &= \sum'_{\substack{n = h^b r^a \\ ah/b \leq r}} h^{-\frac{b+2a}{2(a+b)}} r^{-\frac{2b+a}{2(a+b)}}. \end{aligned}$$

It is easy to check that if that if $h^b r^a \leq \min(H^{a+b}, T^{a/b}) \mathcal{L}^{-a-a^2/b-1}$, $ah/b \leq r$, then $h \leq H, r \leq n_{J+1, h}(b, a)$. Thus

$$(3.10) \quad g^*(b, a; n, H, J) = g(b, a; n), \quad n \leq \min(H^{a+b}, T^{a/b}) \mathcal{L}^{-a-a^2/b-1}.$$

Suppose z is a parameter such that $T^\varepsilon \ll z \leq \min(H^{a+b}, T^{a/b}) \mathcal{L}^{-b-b^2/a-1}$ and define

$$\begin{aligned} R_{11}^*(a, b; x) : &= \frac{c_1(a, b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq z} g(a, b; n) \\ &\quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right) \\ R_{12}^*(a, b; x) : &= R_1^*(a, b; x) - R_{11}^*(a, b; x), \\ R_{11}^*(b, a; x) : &= \frac{c_1(b, a)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq z} g(b, a; n) \\ &\quad \times \cos \left(2\pi c_2(b, a) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right) \\ R_{12}^*(b, a; x) : &= R_1^*(b, a; x) - R_{11}^*(b, a; x). \end{aligned}$$

Recalling (3.4) we have

$$\begin{aligned} R_{11}^*(a, b; x) + R_{11}^*(b, a; x) &= \frac{c_1(a, b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq z} (g(a, b; n) + g(b, a; n)) \\ &\quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right). \end{aligned}$$

From the definition of $g(a, b; n)$ and $g(b, a; n)$ we have

$$\begin{aligned}
& g(a, b; n) + g(b, a; n) \\
&= \sum'_{\substack{n=h^a r^b \\ bh/a \leq r}} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} + \sum'_{\substack{n=h^b r^a \\ ah/b \leq r}} h^{-\frac{b+2a}{2(a+b)}} r^{-\frac{2b+a}{2(a+b)}} \\
&= \sum'_{\substack{n=h^a r^b \\ bh/a \leq r}} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} + \sum'_{\substack{n=r^b h^a \\ ar/b \leq h}} r^{-\frac{b+2a}{2(a+b)}} h^{-\frac{2b+a}{2(a+b)}} \\
&= \sum_{n=h^a r^b} h^{-\frac{a+2b}{2(a+b)}} r^{-\frac{2a+b}{2(a+b)}} = g_{a,b}(n).
\end{aligned}$$

Define

$$\begin{aligned}
\Delta_{a,b}^*(x, z) &:= \frac{c_1(a, b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{1 \leq n \leq z} g_{a,b}(n) \\
&\quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right).
\end{aligned}$$

Combining the above estimates we get

$$\begin{aligned}
(3.11) \quad \Delta_{a,b}(x) &= \Delta_{a,b}^*(x, z) + E_{a,b}(x), \\
E_{a,b}(x) : &= R_{12}^*(a, b; x) + R_{12}^*(b, a; x) + R_2(a, b; x) \\
&\quad + R_2(b, a; x) + O(\mathcal{L}^2).
\end{aligned}$$

The formula (3.11) can be viewed as a truncated Voronoi's formula.

4 On the series $\sum_{n=1}^{\infty} g_{a,b}^2(n)$

In this section we shall prove that the infinite series $\sum_{n=1}^{\infty} g_{a,b}^2(n)$ is convergent. Without loss of generality, we suppose $a < b$. By the definition of $g_{a,b}(n)$ it is easy to see that

$$(4.1) \quad g_{a,b}(n) = n^{-\frac{2a+b}{(2a+2b)b}} \sum_{n=h^a r^b} h^{-\frac{b-a}{a}} = n^{-\frac{2a+b}{(2a+2b)b}} g_{a,b}^*(n),$$

say, where

$$g_{a,b}^*(n) := \sum_{n=h^a r^b} h^{-\frac{b-a}{a}}$$

is a multiplicative function. So we have

$$(4.2) \quad \sum_{n=1}^{\infty} \frac{g_{a,b}^{*2}(n)}{n^s} = \prod_p \left(1 + \sum_{\alpha=1}^{\infty} \frac{g_{a,b}^{*2}(p^\alpha)}{p^{\alpha s}} \right) \quad (\Re s > 1).$$

We shall show that

$$(4.3) \quad \sum_{n=1}^{\infty} \frac{g_{a,b}^{*2}(n)}{n^s} = \zeta(bs) \zeta(as + \frac{2(b-a)}{a}) G_{a,b}(s),$$

where $G_{a,b}(s)$ is analytic for $\Re s > (3a - 2b)/a^2$.

We consider two cases .

Case 1. $a = 1$

If $1 \leq \alpha \leq b-1$, then $p^\alpha = h^a r^b$ implies $h = p^\alpha, r = 1$. So

$$g_{a,b}^*(p^\alpha) = p^{-\alpha(b-1)}.$$

Suppose $\alpha = mb$ with $m \geq 1$. Then $p^\alpha = h^a r^b$ implies $h = p^{jb}, r = p^{(m-j)b}, j = 0, 1, 2, \dots, m$. So

$$g_{a,b}^*(p^\alpha) = \sum_{j=0}^m p^{-jb(b-1)}.$$

Suppose $\alpha = mb + u$ with $m \geq 1$ and $1 \leq u \leq b-1$. Then $p^\alpha = h^a r^b$ implies $h = p^{jb+u}, r = p^{(m-j)b}, j = 0, 1, 2, \dots, m$. So

$$g_{a,b}^*(p^\alpha) = \sum_{j=0}^m p^{-(jb+u)(b-1)} = p^{-u(b-1)} + p^{-u(b-1)} \sum_{j=1}^m p^{-jb(b-1)}.$$

From the above we get ($p \geq 2, \Re s = \sigma > 1$)

$$\begin{aligned} & 1 + \sum_{\alpha=1}^{\infty} \frac{g_{a,b}^{*2}(p^\alpha)}{p^{\alpha s}} \\ &= 1 + \sum_{\alpha=1}^{b-1} \frac{p^{-2\alpha(b-1)}}{p^{\alpha s}} + \sum_{m=1}^{\infty} \frac{(\sum_{j=0}^m p^{-jb(b-1)})^2}{p^{mbs}} + \sum_{m=1}^{\infty} \sum_{u=1}^{b-1} \frac{(\sum_{j=0}^m p^{-(jb+u)(b-1)})^2}{p^{(mb+u)s}} \\ &= 1 + p^{-bs} + O(p^{-\sigma-2(b-1)}). \end{aligned}$$

Inserting this formula into (4.2) we get that (4.3) is true for $a = 1$.

Case 2. $a \geq 2$

Let $k = [b/a]$. Then $(a, b) = 1$ implies that $ka < b < ka + a$.

Consider the equation

$$(4.4) \quad ua + vb = \alpha, \quad u, v \in \mathbb{Z}.$$

Let $v(\alpha; a, b)$ denote the number of non-negative solutions of the equation (4.4). If $v(\alpha; a, b) = 0$, then $g_{a,b}^*(p^\alpha) = 0$. Especially, it is easy to see that if $\alpha \leq b - 1, a \nmid \alpha$, then $v(\alpha; a, b) = 0$. Hence

$$g_{a,b}^*(p^\alpha) = 0, \quad (\alpha \leq b - 1, a \nmid \alpha).$$

If $\alpha = ja, (j = 1, 2, \dots, k)$, then the equation (4.4) has only one solution $(u, v) = (j, 0)$. So $p^\alpha = h^a r^b$ implies that $h = p^j, r = 1$, and hence

$$g_{a,b}^*(p^\alpha) = p^{-\frac{j(b-a)}{a}}.$$

Later suppose $\alpha \geq b$ and $v(\alpha; a, b) > 0$. Let (u_0, v_0) denote a special non-negative solution of (4.4) such that $0 \leq u_0 < b$. Then all non-negative solutions of (4.4) are $(u, v) = (u_0 + bt, v_0 - at), 0 \leq t \leq v(\alpha; a, b)$. Hence we have

$$g_{a,b}^*(p^\alpha) = \sum_{j=0}^{v(\alpha; a, b)-1} p^{-\frac{(u_0+bj)(b-a)}{a}}.$$

Especially when $b|\alpha$, $(0, \alpha/b)$ is a non-negative solution of (4.4). Thus

$$g_{a,b}^*(p^\alpha) = \sum_{j=0}^{v(\alpha; a, b)-1} p^{-\frac{bj(b-a)}{a}}, \quad \alpha \equiv 0 \pmod{b}.$$

From the above we get

$$\begin{aligned} & 1 + \sum_{\alpha=1}^{\infty} \frac{g_{a,b}^{*2}(p^\alpha)}{p^{\alpha s}} \\ &= 1 + \sum_{j=1}^k \frac{p^{-2j(b-a)/a}}{p^{jas}} + \sum_{\alpha > b, b|\alpha} \frac{g_{a,b}^{*2}(p^\alpha)}{p^{\alpha s}} + \sum_{\alpha > b, b \nmid \alpha} \frac{g_{a,b}^{*2}(p^\alpha)}{p^{\alpha s}} \\ &= 1 + p^{-bs} + O(p^{-a\sigma - \frac{2(b-a)}{a}}). \end{aligned}$$

Inserting this formula into (4.2) we get that (4.3) is true in this case.

It is easy to check that $\zeta(as + \frac{2b-2a}{a})$ has a simple pole at $s = (3a - 2b)/a^2$. Since $b > a \geq 1$, it is easy to see that $(3a - 2b)/a^2 < 1/b$. From (4.3) and Perron's formula we get that for any $X > 2$,

$$\sum_{X < n \leq 2X} g_{a,b}^{*2}(n) \ll X^{1/b}.$$

Hence we get

$$(4.5) \quad \sum_{X < n \leq 2X} g_{a,b}^2(n) \ll X^{\frac{1}{b} - \frac{2a+b}{(a+b)b}} \ll X^{-\frac{a}{(a+b)b}},$$

which implies the convergence of the infinite series $\sum_{n=1}^{\infty} g_{a,b}^2(n)$. From (4.5) we also get

$$(4.6) \quad \sum_{n \leq X} g_{a,b}^2(n) = \sum_{n=1}^{\infty} g_{a,b}^2(n) + O(X^{-\frac{a}{(a+b)b}})$$

and

$$(4.7) \quad \sum_{n > X} g_{a,b}^2(n) = O(X^{-\frac{a}{(a+b)b}}).$$

5 On a special sum

Suppose a, b are fixed natural integers, T is a large parameter. Define

$$S_{a,b}(T) : \quad = \sum_2 (h_1 h_2)^{-\frac{2b+a}{2a+2b}} (r_1 r_2)^{-\frac{2a+b}{2a+2b}} \\ \times \min \left(T^{\frac{1}{a+b}}, \frac{1}{\left| h_1^{\frac{a}{a+b}} r_1^{\frac{b}{a+b}} - h_2^{\frac{a}{a+b}} r_2^{\frac{b}{a+b}} \right|} \right),$$

where

$$SC(\Sigma_2) : \quad h_1^a r_1^b \leq T^{100(a+b)}, h_2^a r_2^b \leq T^{100(a+b)}, \\ 0 < |h_1^{\frac{a}{a+b}} r_1^{\frac{b}{a+b}} - h_2^{\frac{a}{a+b}} r_2^{\frac{b}{a+b}}| < \frac{1}{10} h_1^{\frac{a}{2a+2b}} r_1^{\frac{b}{2a+2b}} h_2^{\frac{a}{2a+2b}} r_2^{\frac{b}{2a+2b}}.$$

In this section we shall estimate the sum $S_{a,b}(T)$, which is very important in our proof.

5.1 On a Diophantine inequality

Suppose α and β are fixed non-zero real numbers, $H_1 \geq 1, H_2 \geq 1, R_1 \geq 1, R_2 \geq 1$ are large real numbers, $\delta > 0$. Let $\mathcal{A}(H_1, H_2, R_1, R_2; \delta)$ denote the number of the solutions of the inequality

$$(5.1) \quad \left| h_1^\alpha r_1^\beta - h_2^\alpha r_2^\beta \right| \leq \delta, h_1 \sim H_1, h_2 \sim H_2, r_1 \sim R_1, r_2 \sim R_2.$$

Lemma 5.1. We have

$$\begin{aligned} \mathcal{A}(H_1, H_2, R_1, R_2; \delta) &\ll \delta(H_1 H_2)^{1-\alpha/2} (R_1 R_2)^{1-\beta/2} \\ &\quad + (H_1 H_2 R_1 R_2)^{1/2} (\log 2H_1 H_2 R_1 R_2)^2, \end{aligned}$$

where the implied constant depends only on α, β .

Remark. When $H_1 = H_2, R_1 = R_2$, Lemma 5.1 reduces to Lemma 1 of Fouvry and Iwaniec[4]. Here Lemma 5.1 is more general.

Proof. We follow the proof of Lemma 1 of Fouvry and Iwaniec [4]. Suppose u and v are two positive integers and let $\mathcal{A}_{u,v}(H_1, H_2, R_1, R_2; \delta)$ denote the number of solutions of the inequality (5.1) with $(r_1, r_2) = u, (h_1, h_2) = v$. Set $r_j = m_j u, h_j = l_j v (j = 1, 2)$, then (m_1, m_2, l_1, l_2) satisfies

$$(5.2) \quad \left| \frac{m_1^\beta}{m_2^\beta} - \frac{l_2^\alpha}{l_1^\alpha} \right| \leq c(\alpha)c(\beta)\delta R_2^{-\beta} H_1^{-\alpha},$$

$$(5.3) \quad \left| \frac{m_2^\beta}{m_1^\beta} - \frac{l_1^\alpha}{l_2^\alpha} \right| \leq c(\alpha)c(\beta)\delta R_1^{-\beta} H_2^{-\alpha},$$

where

$$c(\gamma) = \begin{cases} 1, & \text{if } \gamma > 0, \\ 2^{-\gamma}, & \text{if } \gamma < 0. \end{cases}$$

It is easy to see that $\frac{m_1^\beta}{m_2^\beta}$ is $c_1(\beta)u^2(R_1/R_2)^{\beta-1}R_2^{-2}$ -spaced, where $c_1(\beta) > 0$ is some positive constant. So from (5.2) we get

$$\begin{aligned} \mathcal{A}_{u,v}(H_1, H_2, R_1, R_2; \delta) &\ll \frac{H_1 H_2}{v^2} \left(1 + \frac{\delta R_2^{-\beta} H_1^{-\alpha}}{u^2 (R_1/R_2)^{\beta-1} R_2^{-2}} \right) \\ &\ll \frac{H_1 H_2}{v^2} + \frac{\delta H_1 H_2 R_1 R_2}{u^2 v^2 R_1^\beta H_1^\alpha}. \end{aligned}$$

Similarly, $\frac{m_2^\beta}{m_1^\beta}$ is $c_2(\beta)u^2(R_2/R_1)^{\beta-1}R_1^{-2}$ -spaced for some positive constant $c_2(\beta)$, so from (5.3) we get

$$\begin{aligned} \mathcal{A}_{u,v}(H_1, H_2, R_1, R_2; \delta) &\ll \frac{H_1 H_2}{v^2} \left(1 + \frac{\delta R_1^{-\beta} H_2^{-\alpha}}{u^2 (R_2/R_1)^{\beta-1} R_1^{-2}} \right) \\ &\ll \frac{H_1 H_2}{v^2} + \frac{\delta H_1 H_2 R_1 R_2}{u^2 v^2 R_2^\beta H_2^\alpha}. \end{aligned}$$

From the above two estimates we get

$$\begin{aligned} (5.4) \quad \mathcal{A}_{u,v}(H_1, H_2, R_1, R_2; \delta) &\ll \frac{H_1 H_2}{v^2} + \frac{\delta H_1 H_2 R_1 R_2}{u^2 v^2} \min(H_1^{-\alpha} R_1^{-\beta}, H_2^{-\alpha} R_2^{-\beta}) \\ &\ll \frac{H_1 H_2}{v^2} + \frac{\delta (H_1 H_2)^{1-\alpha/2} (R_1 R_2)^{1-\beta}}{u^2 v^2}, \end{aligned}$$

if we note that $\min(x, y) \leq x^{1/2}y^{1/2}$ for $x > 0, y > 0$.

Similarly we have

$$\mathcal{A}_{u,v}(H_1, H_2, R_1, R_2; \delta) \ll \frac{R_1 R_2}{u^2} + \frac{\delta(H_1 H_2)^{1-\alpha/2} (R_1 R_2)^{1-\beta}}{u^2 v^2},$$

which combining (5.4) gives

$$\begin{aligned} & \mathcal{A}_{u,v}(D_1, D_2, N_1, N_2; \delta) \\ & \ll \frac{\delta(H_1 H_2)^{1-\alpha/2} (R_1 R_2)^{1-\beta}}{u^2 v^2} + \min\left(\frac{R_1 R_2}{u^2}, \frac{H_1 H_2}{v^2}\right) \\ & \ll \frac{\delta(H_1 H_2)^{1-\alpha/2} (R_1 R_2)^{1-\beta}}{u^2 v^2} + \frac{(H_1 H_2 R_1 R_2)^{1/2}}{uv}. \end{aligned}$$

Summing over u and v completes the proof of Lemma 5.1. $\square$

5.2 Estimate of the sum $S_{a,b}(T)$

In this subsection we estimate the sum $S_{a,b}(T)$. For simplicity, let

$$\eta := h_1^{\frac{a}{a+b}} r_1^{\frac{b}{a+b}} - h_2^{\frac{a}{a+b}} r_2^{\frac{b}{a+b}}.$$

By a splitting argument we get for some

$$(1, 1, 1, 1) \ll (H_1, H_2, R_1, R_2) \ll (T^{\frac{100(a+b)}{a}}, T^{\frac{100(a+b)}{a}}, T^{\frac{100(a+b)}{b}}, T^{\frac{100(a+b)}{b}})$$

that

$$(5.5) \quad S_{a,b}(T) \ll \mathcal{L}^4 U_{a,b}(T; H_1, H_2, R_1, R_2),$$

where

$$\begin{aligned} U_{a,b}(T; H_1, H_2, R_1, R_2) &:= \sum_3 \frac{1}{(h_1 h_2)^{\frac{a+2b}{2a+2b}} (r_1 r_2)^{\frac{b+2a}{2a+2b}}} \min(T^{\frac{1}{a+b}}, |\eta|^{-1}), \\ SC(\Sigma_3) : h_j &\sim H_j, r_j \sim R_j, (j = 1, 2), |\eta| < \frac{1}{10} h_1^{\frac{a}{2a+2b}} r_1^{\frac{b}{2a+2b}} h_2^{\frac{a}{2a+2b}} r_2^{\frac{b}{2a+2b}}. \end{aligned}$$

By Lemma 5.1 with $(\alpha, \beta) = (a/(a+b), b/(a+b))$, the contribution of $T^{1/(a+b)}$ is

$$\begin{aligned} & \ll \frac{T^{\frac{1}{a+b}}}{(H_1 H_2)^{\frac{a+2b}{2a+2b}} (R_1 R_2)^{\frac{b+2a}{2a+2b}}} \times \mathcal{A}(H_1, H_2, R_1, R_2; T^{-\frac{1}{a+b}}) \\ & \ll 1 + \frac{T^{\frac{1}{a+b}}}{(H_1 H_2)^{\frac{b}{2a+2b}} (R_1 R_2)^{\frac{a}{2a+2b}}} \mathcal{L}^2. \end{aligned}$$

The condition $SC(\Sigma_3)$ implies that $h_1^a r_1^b \asymp h_2^a r_2^b$. So by the mean value theorem we get for some $x_0 \asymp h_1^a r_1^b$ that

$$(5.6) \quad |\eta| = \frac{1}{a+b} x_0^{\frac{1}{a+b}-1} |h_1^a r_1^b - h_2^a r_2^b| \gg (h_1^a r_1^b)^{\frac{1}{a+b}-1},$$

which combining the inequality $|\eta| \leq T^{-1/(a+b)}$ gives

$$H_1^a R_1^b \gg T^{1/(a+b-1)}.$$

Similarly we have

$$H_2^a R_2^b \gg T^{1/(a+b-1)}.$$

If $a \leq b$, then combining the above estimates we get that the contribution of $T^{1/(a+b)}$ is

$$\begin{aligned} & \ll 1 + \frac{T^{\frac{1}{a+b}}}{(H_1^a R_1^b)^{\frac{a}{(2a+2b)b}} (H_2^a R_2^b)^{\frac{a}{(2a+2b)b}} (H_1 H_2)^{\frac{b-a}{2b}}} \mathcal{L}^2 \\ & \ll T^{\frac{1}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^2. \end{aligned}$$

If $a > b$, then the contribution of $T^{1/(a+b)}$ is

$$\begin{aligned} & \ll 1 + \frac{T^{\frac{1}{a+b}}}{(H_1^a R_1^b)^{\frac{b}{(2a+2b)a}} (H_2^a R_2^b)^{\frac{b}{(2a+2b)a}} (R_1 R_2)^{\frac{a-b}{2a}}} \mathcal{L}^2 \\ & \ll T^{\frac{1}{a+b} - \frac{b}{a(a+b)(a+b-1)}} \mathcal{L}^2. \end{aligned}$$

Namely, the the contribution of $T^{1/(a+b)}$ is

$$(5.7) \quad \ll T^{\frac{1}{a+b} - \frac{1}{(a+b)(a+b-1)} \times \frac{\min(a,b)}{\max(a,b)}} \mathcal{L}^2.$$

Now we consider the contribution of $1/|\eta|$. Suppose first that $a \leq b$. By a splitting argument the contribution of $1/|\eta|$ is

$$\begin{aligned} & \ll \frac{\mathcal{L}}{(H_1 H_2)^{\frac{a+2b}{2a+2b}} (R_1 R_2)^{\frac{b+2a}{2a+2b}} \delta} \times \mathcal{A}(H_1, H_2, R_1, R_2; \delta) \\ & \ll \mathcal{L} + \frac{1}{(H_1 H_2)^{\frac{b}{2a+2b}} (R_1 R_2)^{\frac{a}{2a+2b}} \delta} \mathcal{L}^3 \end{aligned}$$

for some $T^{-1/(a+b)} \ll \delta \ll H_1^{\frac{a}{2a+2b}} R_1^{\frac{b}{2a+2b}} H_2^{\frac{a}{2a+2b}} R_2^{\frac{b}{2a+2b}}$. From (5.6) we get

$$\delta \gg (H_1^a R_1^b)^{\frac{1}{a+b}-1},$$

which combining $T^{-1/(a+b)} \ll \delta$ gives

$$\delta^{-1} \ll \min \left(T^{\frac{1}{a+b}}, (H_1^a R_1^b)^{\frac{a+b-1}{a+b}} \right).$$

Thus the contribution of $1/|\eta|$ is

$$\begin{aligned} & \ll \mathcal{L} + \frac{1}{(H_1^a R_1^b)^{\frac{a}{(2a+2b)b}} (H_2^a R_2^b)^{\frac{a}{(2a+2b)b}} (H_1 H_2)^{\frac{b-a}{2b}} \delta} \mathcal{L}^3 \\ & \ll \mathcal{L} + \frac{1}{(H_1^a R_1^b)^{\frac{a}{(a+b)b}} (H_1 H_2)^{\frac{b-a}{2b}} \delta} \mathcal{L}^3 \\ & \ll \mathcal{L} + \frac{\min \left(T^{\frac{1}{a+b}}, (H_1^a R_1^b)^{\frac{a+b-1}{a+b}} \right)}{(H_1^a R_1^b)^{\frac{a}{(a+b)b}} (H_1 H_2)^{\frac{b-a}{2b}}} \mathcal{L}^3 \\ & \ll \mathcal{L} + (H_1 H_2)^{-\frac{b-a}{2b}} \\ & \quad \times \min \left(T^{\frac{1}{a+b}} (H_1^a R_1^b)^{-\frac{a}{(a+b)b}}, (H_1^a R_1^b)^{\frac{a+b-1}{a+b} - \frac{a}{b(a+b)}} \right) \mathcal{L}^3 \\ & \ll \left(T^{\frac{1}{a+b}} (H_1^a R_1^b)^{-\frac{a}{(a+b)b}} \right)^{1 - \frac{a}{b(a+b-1)}} \\ & \quad \times \left((H_1^a R_1^b)^{\frac{a+b-1}{a+b} - \frac{a}{b(a+b)}} \right)^{\frac{a}{b(a+b-1)}} \mathcal{L}^3 \\ & \ll T^{\frac{1}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^3, \end{aligned}$$

where in the second step we used the fact $H_1^a R_1^b \asymp H_2^a R_2^b$.

Similarly if $a > b$, we can get that the contribution of $1/|\eta|$ is

$$\ll T^{\frac{1}{a+b} - \frac{b}{a(a+b)(a+b-1)}} \mathcal{L}^3.$$

Hence combining the above we see that the contribution of $1/|\eta|$ is

$$(5.8) \quad \ll T^{\frac{1}{a+b} - \frac{1}{(a+b)(a+b-1)} \times \frac{\min(a,b)}{\max(a,b)}} \mathcal{L}^3.$$

From (5.5), (5.7) and (5.8) we get the following

Lemma 5.2. Suppose a and b are fixed natural numbers, then we have

$$S_{a,b}(T) \ll T^{\frac{1}{a+b} - \frac{1}{(a+b)(a+b-1)} \times \frac{\min(a,b)}{\max(a,b)}} \mathcal{L}^7.$$

6 Proof of Theorem

In this section we shall prove our Theorem . It suffices to evaluate the integral $\int_T^{2T} \Delta_{a,b}^2(x) dx$ for $T \geq 10$. We always suppose $a < b$.

6.1 Mean square of $\Delta_{a,b}^*(x, z)$

Suppose H and z are parameters such that

$$T^\varepsilon \ll H \ll T^{100(a+b)}, \quad T^\varepsilon \ll z \leq \min(H^{a+b}, T^{a/b}) \mathcal{L}^{-b-b^2/a-1}.$$

In this subsection we study the mean square of $\Delta_{a,b}^*(x, z)$.

By the elementary formula

$$(6.1) \quad \cos u \cos v = \frac{1}{2}(\cos(u-v) + \cos(u+v))$$

we may write

$$(6.2) \quad \begin{aligned} & |\Delta_{a,b}^*(x, z)|^2 \\ &= \frac{c_1^2(a, b)}{\pi^2} x^{\frac{1}{a+b}} \sum_{1 \leq n \leq z} \sum_{1 \leq m \leq z} g_{a,b}(n) g_{a,b}(m) \\ & \quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right) \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} m^{\frac{1}{a+b}} - \frac{\pi}{4} \right) \\ &= S_1(x) + S_2(x) + S_3(x), \end{aligned}$$

where

$$\begin{aligned} S_1(x) &= \frac{c_1^2(a, b)}{2\pi^2} x^{\frac{1}{a+b}} \sum_{1 \leq n \leq z} g_{a,b}^2(n), \\ S_2(x) &= \frac{c_1^2(a, b)}{2\pi^2} x^{\frac{1}{a+b}} \sum_{\substack{n \leq z, m \leq z \\ n \neq m}} g_{a,b}(n) g_{a,b}(m) \\ & \quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} (n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}) \right) \\ S_3(x) &= \frac{c_1^2(a, b)}{2\pi^2} x^{\frac{1}{a+b}} \sum_{n \leq z, m \leq z} g_{a,b}(n) g_{a,b}(m) \\ & \quad \times \sin \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} (n^{\frac{1}{a+b}} + m^{\frac{1}{a+b}}) \right). \end{aligned}$$

By (4.6) we get

$$(6.3) \quad \int_T^{2T} S_1(x) dx = \frac{c_1^2(a, b)}{2\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n) \int_T^{2T} x^{\frac{1}{a+b}} dx + O(T^{\frac{1+a+b}{a+b}} z^{-\frac{a}{(a+b)b}}).$$

By the first derivative test we get

$$\begin{aligned}
\int_T^{2T} S_3(x) dx &\ll \sum_{n \leq z, m \leq z} g_{a,b}(n) g_{a,b}(m) \frac{T}{n^{\frac{1}{a+b}} + m^{\frac{1}{a+b}}} \\
&\ll T \sum_{n \leq z, m \leq z} g_{a,b}(n) g_{a,b}(m) \frac{1}{n^{\frac{1}{2(a+b)}} m^{\frac{1}{2(a+b)}}} \\
&\ll T \left(\sum_{n \leq z} \frac{g_{a,b}(n)}{n^{\frac{1}{2(a+b)}}} \right)^2,
\end{aligned}$$

where in the second step we used the well-known inequality $\alpha^2 + \beta^2 \geq 2\alpha\beta$. By Euler's product we have for $\Re s > 1$ that

$$\sum_{n=1}^{\infty} \frac{g_{a,b}(n)}{n^s} = \zeta\left(as + \frac{a+2b}{2a+2b}\right) \zeta\left(bs + \frac{2a+b}{2a+2b}\right),$$

which can be continued meromorphically to the whole complex plane and has a double pole at $s = 1/2(a+b)$. Thus Perron's formula implies that

$$\sum_{n \leq X} g_{a,b}(n) \ll X^{1/2(a+b)} \log X$$

for any $X > 2$, namely,

$$(6.4) \quad \sum_{n \leq X} g_{a,b}(n) n^{-1/2(a+b)} \ll \log^2 X.$$

From the above we get

$$(6.5) \quad \int_T^{2T} S_3(x) dx \ll T \mathcal{L}^4.$$

Now we consider the contribution of $S_2(x)$. Write

$$(6.6) \quad S_2(x) = S_{21}(x) + S_{22}(x),$$

where

$$\begin{aligned}
S_{2j}(x) &= \frac{c_1^2(a,b)}{2\pi^2} x^{\frac{1}{(a+b)}} \sum_{2j} g_{a,b}(n) g_{a,b}(m) \\
&\quad \times \cos\left(2\pi c_2(a,b) x^{\frac{1}{a+b}} (n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}})\right) \quad (j = 1, 2),
\end{aligned}$$

$$SC(\Sigma_{21}) : \quad n, m \leq z, |n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}| \geq \frac{(nm)^{1/(2a+2b)}}{10},$$

$$SC(\Sigma_{22}) : \quad n, m \leq z, 0 < |n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}| < \frac{(nm)^{1/(2a+2b)}}{10}.$$

Similar to the case $S_3(x)$, we have

$$\int_T^{2t} S_{21}(x)dx \ll T\mathcal{L}^4.$$

By the first derivative test and Lemma 5.2 we get

$$\begin{aligned} \int_T^{2T} S_{22}(x)dx &\ll T \sum_{22} g_{a,b}(n)g_{a,b}(m) \min \left(T^{\frac{1}{a+b}}, \frac{1}{|n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}|} \right) \\ &\ll TS_{a,b}(T) \ll T^{1+\frac{1}{a+b}-\frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7. \end{aligned}$$

From the above two estimates we get

$$(6.7) \quad \int_T^{2T} S_2(x)dx \ll T^{1+\frac{1}{a+b}-\frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7.$$

From (6.2), (6.3), (6.5) and (6.7) we have

$$\begin{aligned} (6.8) \quad \int_T^{2T} |\Delta_{a,b}^*(x, z)|^2 dx &= \frac{c_1^2(a, b)}{2\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n) \int_T^{2T} x^{\frac{1}{a+b}} dx \\ &\quad + O(T^{\frac{1+a+b}{a+b}} z^{-\frac{a}{(a+b)b}} + T^{\frac{1+a+b}{a+b}-\frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7). \end{aligned}$$

6.2 Mean squares of $R_{12}^*(a, b; x)$ and $R_{12}^*(b, a; x)$

In this subsection we shall study the mean squares of $R_{12}^*(a, b; x)$ and $R_{12}^*(b, a; x)$. Recall that

$$\begin{aligned} R_{12}^*(a, b; x) &= \frac{c_1(a, b)}{\pi} x^{\frac{1}{2(a+b)}} \sum_{z \leq n \leq H^a n_{J+1, H}(a, b)} g(a, b; n, H, J) \\ &\quad \times \cos \left(2\pi c_2(a, b) x^{\frac{1}{a+b}} n^{\frac{1}{a+b}} - \frac{\pi}{4} \right). \end{aligned}$$

Thus

$$\begin{aligned} &|R_{12}^*(a, b; x)|^2 \\ &\ll x^{\frac{1}{a+b}} \sum_{z \leq n, m \leq H^a n_{J+1, H}(a, b)} g(a, b; n, H, J) g(a, b; m, H, J) \\ &\quad \times e \left(c_2(a, b) x^{\frac{1}{a+b}} (n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}) \right) \\ &= S_4(x) + S_5(x), \end{aligned}$$

where

$$\begin{aligned}
S_4(x) &= x^{\frac{1}{a+b}} \sum_{z \leq n \leq H^a n_{J+1, H}(a, b)} g^2(a, b; n, H, J), \\
S_5(x) &= x^{\frac{1}{a+b}} \sum_{\substack{z \leq n, m \leq H^a n_{J+1, H}(a, b) \\ n \neq m}} g(a, b; n, H, J) g(a, b; m, H, J) \\
&\quad \times e \left(c_2(a, b) x^{\frac{1}{a+b}} (n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}) \right)
\end{aligned}$$

It is easy to see that

$$(6.9) \quad g(a, b; n, H, J) \leq g(a, b; n) \leq g_{a, b}(n).$$

By (4.7) we get

$$(6.10) \quad \int_T^{2T} S_4(x) dx \ll T^{1+\frac{1}{a+b}} z^{-\frac{a}{b(a+b)}}$$

By the first derivative test and (6.9) we get

$$\begin{aligned}
(6.11) \quad \int_T^{2T} S_5(x) dx &\ll T \sum_{\substack{z \leq n, m \leq H^a T \\ n \neq m}} g_{a, b}(n) g_{a, b}(m) \\
&\quad \times \min \left(T^{\frac{1}{a+b}}, \frac{1}{|n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}|} \right) \\
&\ll T(\Sigma_4 + \Sigma_5),
\end{aligned}$$

where

$$\begin{aligned}
\Sigma_4 &:= \sum_4 g_{a, b}(n) g_{a, b}(m) \min \left(T^{\frac{1}{a+b}}, \frac{1}{|n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}|} \right), \\
\Sigma_5 &:= \sum_5 g_{a, b}(n) g_{a, b}(m) \min \left(T^{\frac{1}{a+b}}, \frac{1}{|n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}|} \right), \\
SC(\Sigma_4) &: z < n, m \leq H^a T, 0 < |n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}| \leq \frac{1}{10} n^{\frac{1}{2a+2b}} m^{\frac{1}{2a+2b}}, \\
SC(\Sigma_5) &: z < n, m \leq H^a T, |n^{\frac{1}{a+b}} - m^{\frac{1}{a+b}}| \geq \frac{1}{10} n^{\frac{1}{2a+2b}} m^{\frac{1}{2a+2b}}.
\end{aligned}$$

By Lemma 5.2 we have

$$(6.12) \quad \Sigma_4 \ll S_{a, b}(T) \ll T^{\frac{1}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7.$$

For Σ_5 by (6.4) we get

$$(6.13) \quad \Sigma_5 \ll \left(\sum_{n \leq H^a T} g_{a,b}(n) n^{-1/(2a+2b)} \right)^2 \ll \log^4 X.$$

From (6.10)-(6.13) we have

$$(6.14) \quad \int_T^{2T} |R_{12}^*(a, b; x)|^2 dx \ll T^{1+\frac{1}{a+b}} z^{-\frac{a}{b(a+b)}} + T^{\frac{1+a+b}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7.$$

Similarly, we have

$$(6.15) \quad \int_T^{2T} |R_{12}^*(b, a; x)|^2 dx \ll T^{1+\frac{1}{a+b}} z^{-\frac{a}{b(a+b)}} + T^{\frac{1+a+b}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7.$$

6.3 Mean squares of $R_2(a, b; x)$ and $R_2(b, a; x)$

In this subsection we shall study the mean squares of $R_2(a, b; x)$ and $R_2(b, a; x)$. Recall that

$$R_2(a, b; x) = O \left(\sum_{m \leq x^{1/(a+b)}} \min \left(1, \frac{1}{H \left\| \frac{x^{1/a}}{m^{b/a}} \right\|} \right) \right).$$

So we have

$$\begin{aligned}
& \int_T^{2T} R_2(a, b; x) dx \\
& \ll \sum_{m \leq 2T^{1/(a+b)}} \int_T^{2T} \min \left(1, \frac{1}{H \left\| \frac{x^{1/a}}{m^{b/a}} \right\|} \right) dx \\
& \ll \sum_{m \leq 2T^{1/(a+b)}} \int_{T^{1/a}/m^{b/a}}^{(2T)^{1/a}/m^{b/a}} \min \left(1, \frac{1}{H \|u\|} \right) m^b u^{a-1} du \\
& \ll \sum_{m \leq 2T^{1/(a+b)}} m^b \left(\frac{T^{1/a}}{m^{b/a}} \right)^{a-1} \int_{T^{1/a}/m^{b/a}}^{(2T)^{1/a}/m^{b/a}} \min \left(1, \frac{1}{H \|u\|} \right) du \\
& \ll \sum_{m \leq 2T^{1/(a+b)}} m^b \left(\frac{T^{1/a}}{m^{b/a}} \right)^{a-1} \frac{T^{1/a}}{m^{b/a}} \int_0^1 \min \left(1, \frac{1}{H \|u\|} \right) du \\
& \ll T \sum_{m \leq 2T^{1/(a+b)}} \int_0^1 \min \left(1, \frac{1}{H \|u\|} \right) du \\
& \ll T^{1+1/(a+b)} \int_0^{1/2} \min \left(1, \frac{1}{H \|u\|} \right) du \\
& \ll T^{1+1/(a+b)} \left(\int_0^{1/H} du + \int_{1/H}^{1/2} \frac{1}{Hu} du \right) \\
& \ll T^{1+1/(a+b)} H^{-1} \mathcal{L},
\end{aligned}$$

which combining the trivial bound $R_2(a, b; x) \ll T^{1/(a+b)}$ gives

$$(6.16) \quad \int_T^{2T} R_2^2(a, b; x) dx \ll T^{1+2/(a+b)} H^{-1} \mathcal{L}$$

Similarly we have

$$(6.17) \quad \int_T^{2T} R_2^2(b, a; x) dx \ll T^{1+2/(a+b)} H^{-1} \mathcal{L}.$$

6.4 Completion of the proof of Theorem

In this subsection we complete the proof of Theorem . We take $H = T^{10(a+b)}$ and $z = T^{a/b} \mathcal{L}^{-b-b^2/a-1}$. From (3.11), (6.11)-(6.14) we get

$$(6.18) \quad \int_T^{2T} E_{a,b}^2(x) dx \ll T^{\frac{1+a+b}{a+b} - \frac{a}{b(a+b)(a+b-1)}} \mathcal{L}^7,$$

which combining (6.8) and Cauchy's inequality gives

$$(6.19) \quad \int_T^{2T} E_{a,b}(x) \Delta_{a,b}^*(x, z) dx \ll T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \mathcal{L}^{7/2}.$$

From (6.8), (6.18) and (6.19) we get

$$(6.20) \quad \int_T^{2T} \Delta_{a,b}^2(x) dx = \frac{c_1^2(a, b)}{2\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n) \int_T^{2T} x^{\frac{1}{a+b}} dx \\ + O(T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \mathcal{L}^{7/2}).$$

From (6.20) and a splitting argument we get

$$(6.21) \quad \int_1^T \Delta_{a,b}^2(x) dx = \frac{c_1^2(a, b)}{2\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n) \int_1^T x^{\frac{1}{a+b}} dx \\ + O(T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \mathcal{L}^{7/2}) \\ = \frac{c_1^2(a, b)(a+b)}{2(1+a+b)\pi^2} \sum_{n=1}^{\infty} g_{a,b}^2(n) T^{\frac{1+a+b}{a+b}} \\ + O(T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \mathcal{L}^{7/2}) \\ = c_{a,b} T^{\frac{1+a+b}{a+b}} + O(T^{\frac{1+a+b}{a+b} - \frac{a}{2b(a+b)(a+b-1)}} \mathcal{L}^{7/2}).$$

References

- [1] X. D. Cao, On the general divisor problem(in Chinese), Acta Mathematica sinica, Vol.**36**(5)(1993), 644-653.
- [2] K. Corrádi and I. Katai, A comment on K. S. Ganggadharan's paper entitled "Two classical lattice point problems", Magyar Tud. Akad. Mat. Fiz. Oszt. Közl. **17**(1967), 89-97.
- [3] H. Cramér, Über zwei Sätze von Herrn G. H. Hardy, Math. Z.**15**(1922), 201-210.
- [4] E. Fouvry and H. Iwaniec, Exponential sums with monomials, J. of Number Theory, Vol. **33**(3)(1989), 311—333 .
- [5] J. L. Hafner, New omega theorems for two classical lattice point problems, Invent. Math. **63**(1981), 181-186.
- [6] J. L. Hafner, New omega results in a weighted divisor problem, J. of Number Theory Vol. **28**(1988), 240-257.

- [7] D. R. Heath-Brown, The distribution and moments of the error term in the Dirichlet divisor problem, *Acta Arith.* **60** (1992), 389-415.
- [8] D. R. Heath-Brown, The Piatetski-Shapiro prime theorem, *J. of Number theory*, Vol. **16**(1983), 242-266.
- [9] M. N. Huxley, Exponential sums and Lattice points III, *Proc. London Math. Soc.*, Vol. **87**(3)(2003), 591-609.
- [10] A. Ivić, Lectures on mean values of the Riemann zeta-function, *Lectures On Math. and Physics* 82, Tata Inst. Fund. Res., Bombay, 1991.
- [11] A. Ivić, The Riemann zeta-function. John, Wiley and Sons, 1985.
- [12] A. Ivić, The general divisor problem, *J. of Number Theory* **27**(1987), 73-91.
- [13] A. Ivić and P. Sargos, On the higher power moments of the error term in the divisor problem, *Illinois Journal of Math.* **51**(2)(2007), 353-377.
- [14] E. Krätzel, Ein Teilerproblem, *J. Reine Angew. Math.* **235**(1969), 150-174.
- [15] E. Krätzel, Lattice points, *Deutsch. Verlag Wiss.*, Berlin, 1988.
- [16] M. Kühleitner and W. G. Nowak, The asymptotic behaviour of the mean-square of fractional part sums, *Proc. Edinb. Math. Soc.* 43 (2000), 309-323.
- [17] S. H. Min, The methods of number theory(in Chinese), Science Press, Beijing: 1981.
- [18] E. Preissmann, Sur la moyenne quadratique du terme de reste du problème du cercle, *C. R. Acad. Sci. Paris Sr. I* 306 (1988), 151-154.
- [19] R.A. Rankin, Van Der Corput's method and the theory of exponent pairs, *Quart. J. Math. Oxford* (2)**6**(1955), 147-153.
- [20] H.-E. Richert, Über die Anzahl Gruppen gegebener Ordnung (I), *Math. Z.* **56**(1952), 21-32.
- [21] P. G. Schmidt, Abschätzungen bei unsymmetrischen Gitterpunktproblem, *Dissertation*, Göttingen, 1964.
- [22] K. C. Tong, On divisor problem III, *Acta math. Sinica* **6** (1956), 515-541.
- [23] Kai-Man Tsang, Higher-power moments of $\Delta(x)$, $E(t)$ and $P(x)$, *Proc. London Math. Soc.*(3)**65**(1992), 65-84.

- [24] I. M. Vinogradov, Special variants of the method of trigonometric sums, (Nauka, Moscow), 1976; English transl. in his Selected works (Springer-Verlag), 1985.
- [25] Wenguang Zhai , On higher-power moments of $\Delta(x)$, Acta Arith. Vol.**112**(2004), 1-24.
- [26] Wenguang Zhai, On higher-power moments of $\Delta(x)$ (II), Acta Arith. Vol.**114**(2004), 35-54.
- [27] Wenguang Zhai, On higher-power moments of $\Delta(x)$ (III), Acta Arith. **118** (2005), 263–281.

Authors' addresses:

Wenguang Zhai,
 School of Mathematical Sciences,
 Shandong Normal University,
 Jinan, Shandong, 250014,
 P.R.China
 E-mail: zhaiwg@hotmail.com

Xiaodong Cao
 Dept. of Mathematics and Physics,
 Beijing Institute of Petro-Chemical Technology,
 Beijing, 102617, P.R. China
 Email: caoxiaodong@bipt.edu.cn